

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 487/04, A61K 31/519	A1	(11) International Publication Number: WO 00/53605 (43) International Publication Date: 14 September 2000 (14.09.00)
(21) International Application Number: PCT/US00/05903 (22) International Filing Date: 8 March 2000 (08.03.00) (30) Priority Data: 60/123,902 11 March 1999 (11.03.99) US (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BILODEAU, Mark, T. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). FRALEY, Mark, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). HUNGATE, Randall, W. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: TYROSINE KINASE INHIBITORS (57) Abstract The present invention relates to compounds which inhibit, regulate and/or modulate tyrosine kinase signal transduction, compositions which contain these compounds, and methods of using them to treat tyrosine kinase-dependent diseases and conditions, such as angiogenesis, cancer, tumor growth, atherosclerosis, age related macular degeneration, diabetic retinopathy, inflammatory diseases, and the like in mammals.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

TYROSINE KINASE INHIBITORS

RELATED APPLICATIONS

- 5 This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application 60/123,902, filed March 11, 1999.

BACKGROUND OF THE INVENTION

- 10 The present invention relates to compounds which inhibit, regulate and/or modulate tyrosine kinase signal transduction, compositions which contain these compounds, and methods of using them to treat tyrosine kinase-dependent diseases and conditions, such as angiogenesis, cancer, tumor growth, atherosclerosis, age related macular degeneration, diabetic retinopathy, inflammatory diseases, and the like in mammals.

- 15 Tyrosine kinases are a class of enzymes that catalyze the transfer of the terminal phosphate of adenosine triphosphate to tyrosine residues in protein substrates. Tyrosine kinases are believed, by way of substrate phosphorylation, to play critical roles in signal transduction for a number of cell functions. Though the exact mechanisms of signal transduction is still
20 unclear, tyrosine kinases have been shown to be important contributing factors in cell proliferation, carcinogenesis and cell differentiation.

- Tyrosine kinases can be categorized as receptor type or non-receptor type. Receptor type tyrosine kinases have an extracellular, a transmembrane, and an intracellular portion, while non-receptor type tyrosine
25 kinases are wholly intracellular.

- The receptor-type tyrosine kinases are comprised of a large number of transmembrane receptors with diverse biological activity. In fact, about twenty different subfamilies of receptor-type tyrosine kinases have been identified. One tyrosine kinase subfamily, designated the HER subfamily, is
30 comprised of EGFR, HER2, HER3, and HER4. Ligands of this subfamily of

receptors include epithelial growth factor, TGF- α , amphiregulin, HB-EGF, betacellulin and heregulin. Another subfamily of these receptor-type tyrosine kinases is the insulin subfamily, which includes INS-R, IGF-IR, and IR-R. The PDGF subfamily includes the PDGF- α and β receptors, CSFIR, c-kit and
5 FLK-II. Then there is the FLK family which is comprised of the kinase insert domain receptor (KDR), fetal liver kinase-1 (FLK-1), fetal liver kinase-4 (FLK-4) and the fms-like tyrosine kinase-1 (flt-1). The PDGF and FLK families are usually considered together due to the similarities of the two groups. For a detailed discussion of the receptor-type tyrosine kinases, see Plowman et al.,
10 *DN&P* 7(6):334-339, 1994, which is hereby incorporated by reference.

The non-receptor type of tyrosine kinases is also comprised of numerous subfamilies, including Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack, and LIMK. Each of these subfamilies is further sub-divided into varying receptors. For example, the Src subfamily is one of the largest and
15 includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr, and Yrk. The Src subfamily of enzymes has been linked to oncogenesis. For a more detailed discussion of the non-receptor type of tyrosine kinases, see Bolen *Oncogene*, 8:2025-2031 (1993), which is hereby incorporated by reference.

Both receptor-type and non-receptor type tyrosine kinases are
20 implicated in cellular signaling pathways leading to numerous pathogenic conditions, including cancer, psoriasis and hyperimmune responses.

Several receptor-type tyrosine kinases, and the growth factors that bind thereto, have been suggested to play a role in angiogenesis, although some may promote angiogenesis indirectly (Mustonen and Alitalo, *J. Cell Biol.*
25 129:895-898, 1995). One such receptor-type tyrosine kinase is fetal liver kinase 1 or FLK-1. The human analog of FLK-1 is the kinase insert domain-containing receptor KDR, which is also known as vascular endothelial cell growth factor receptor 2 or VEGFR-2, since it binds VEGF with high affinity. Finally, the murine version of this receptor has also been called NYK (Oelrichs
30 et al., *Oncogene* 8(1):11-15, 1993). VEGF and KDR are a ligand-receptor pair

that play an important role in the proliferation of vascular endothelial cells, and the formation and sprouting of blood vessels, termed vasculogenesis and angiogenesis, respectively.

Angiogenesis is characterized by excessive activity of vascular endothelial growth factor (VEGF). VEGF is actually comprised of a family of
5 ligands (Klagsburn and D'Amore, *Cytokine & Growth Factor Reviews* 7:259-270, 1996). VEGF binds the high affinity membrane-spanning tyrosine kinase receptor KDR and the related fms-like tyrosine kinase-1, also known as Flt-1 or vascular endothelial cell growth factor receptor 1 (VEGFR-1). Cell culture and
10 gene knockout experiments indicate that each receptor contributes to different aspects of angiogenesis. KDR mediates the mitogenic function of VEGF whereas Flt-1 appears to modulate non-mitogenic functions such as those associated with cellular adhesion. Inhibiting KDR thus modulates the level of mitogenic VEGF activity. In fact, tumor growth has been shown to be
15 susceptible to the antiangiogenic effects of VEGF receptor antagonists. (Kim et al., *Nature* 362, pp. 841-844, 1993).

Solid tumors can therefore be treated by tyrosine kinase inhibitors since these tumors depend on angiogenesis for the formation of the blood vessels necessary to support their growth. These solid tumors include
20 histiocytic lymphoma, cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung, including lung adenocarcinoma and small cell lung cancer. Additional examples include cancers in which overexpression or activation of Raf-activating oncogenes (e.g., K-ras, erb-B) is observed. Such cancers include pancreatic and breast carcinoma. Accordingly, inhibitors of
25 these tyrosine kinases are useful for the prevention and treatment of proliferative diseases dependent on these enzymes.

The angiogenic activity of VEGF is not limited to tumors. VEGF accounts for most of the angiogenic activity produced in or near the retina in diabetic retinopathy. This vascular growth in the retina leads to visual
30 degeneration culminating in blindness. Ocular VEGF mRNA and protein are

elevated by conditions such as retinal vein occlusion in primates and decreased pO₂ levels in mice that lead to neovascularization. Intraocular injections of anti-VEGF monoclonal antibodies or VEGF receptor immunofusions inhibit ocular neovascularization in both primate and rodent models. Regardless of the
5 cause of induction of VEGF in human diabetic retinopathy, inhibition of ocular VEGF is useful in treating the disease.

Expression of VEGF is also significantly increased in hypoxic regions of animal and human tumors adjacent to areas of necrosis. VEGF is also upregulated by the expression of the oncogenes ras, raf, src and mutant
10 p53 (all of which are relevant to targeting cancer). Monoclonal anti-VEGF antibodies inhibit the growth of human tumors in nude mice. Although these same tumor cells continue to express VEGF in culture, the antibodies do not diminish their mitotic rate. Thus tumor-derived VEGF does not function as an autocrine mitogenic factor. Therefore, VEGF contributes to tumor growth in
15 vivo by promoting angiogenesis through its paracrine vascular endothelial cell chemotactic and mitogenic activities. These monoclonal antibodies also inhibit the growth of typically less well vascularized human colon cancers in athymic mice and decrease the number of tumors arising from inoculated cells.

Viral expression of a VEGF-binding construct of Flk-1, Flt-1,
20 the mouse KDR receptor homologue, truncated to eliminate the cytoplasmic tyrosine kinase domains but retaining a membrane anchor, virtually abolishes the growth of a transplantable glioblastoma in mice presumably by the dominant negative mechanism of heterodimer formation with membrane spanning endothelial cell VEGF receptors. Embryonic stem cells, which
25 normally grow as solid tumors in nude mice, do not produce detectable tumors if both VEGF alleles are knocked out. Taken together, these data indicate the role of VEGF in the growth of solid tumors. Inhibition of KDR or Flt-1 is implicated in pathological angiogenesis, and these receptors are useful in the treatment of diseases in which angiogenesis is part of the overall pathology,
30 e.g., inflammation, diabetic retinal vascularization, as well as various forms of

cancer since tumor growth is known to be dependent on angiogenesis.


(Weidner et al., N. Engl. J. Med., 324, pp. 1-8, 1991).

Accordingly, the identification of small compounds which specifically inhibit, regulate and/or modulate the signal transduction of tyrosine

5 kinases is desirable and is an object of this invention.

SUMMARY OF THE INVENTION


The present invention relates to compounds that are capable of inhibiting, modulating and/or regulating signal transduction of both receptor-type and
10 non-receptor type tyrosine kinases. One embodiment of the present invention is illustrated by a compound of Formula I, and the pharmaceutically acceptable salts and stereoisomers thereof:



I.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of kinases and are illustrated by a compound of Formula I :



I

or a pharmaceutically acceptable salt or stereoisomer thereof, wherein:

5

X is CH or N;

R₁ and R₃ are independently selected from the group consisting of:

10

- 1) H,
- 2) C₁₋₁₀ alkyl,
- 3) C₂₋₁₀ alkenyl,
- 4) C₂₋₁₀ alkynyl,
- 5) aryl,
- 6) halo,
- 7) OH, and
- 8) heterocyclyl,

15

said alkyl, alkenyl, alkynyl, aryl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

20 R₂ is:

- 1) H,
- 2) C₁₋₆ alkyl,
- 3) aryl,

- 4) OH,
- 5) NO₂,
- 6) NH₂, or
- 7) halogen;

5

R₅ is:

- 1) H,
- 2) C₁₋₆ alkyl,
- 3) OH,
- 4) O-C₁₋₆ alkyl,
- 5) halo,
- 6) NH₂, or
- 7) NO₂;

10

15 R₇ and R₈ are independently selected from the group consisting of:

- 1) H,
- 2) C₁₋₁₀ alkyl,
- 3) COR,
- 4) COOR,
- 5) aryl, and
- 6) heterocyclyl,

20

said alkyl, aryl and heterocyclyl optionally substituted with R₉,

or NR₇R₈ are taken together to form a heterocyclic 5-10 membered

25 saturated or unsaturated ring containing, in addition to the nitrogen atom, one to two additional heteroatoms selected from the group consisting of N, O and S, said ring optionally substituted with one or two substituents selected from R_a;

R₉ is aryl or heterocyclyl,
said aryl and heterocyclyl is optionally substituted with from one
to three members selected from R^a;

R₁₀ is:

- 5 1) H,
 2) C₁₋₆ alkyl,
 3) NR₇R₈,
 4) O-C₁₋₆ alkyl,
 5) aryl, or
10 6) heterocyclyl,

said alkyl, aryl, and heterocyclyl is optionally substituted with one to three
members selected from R^a;

R^a is:

- 15 1) C₁₋₁₀ alkyl,
 2) halogen,
 3) NO₂,
 4) OR,
 5) NR₇R₈,
20 6) CN,
 7) aryl, or
 8) heterocyclyl; and

R is H or C₁₋₆ alkyl.
25

A second embodiment is a compound of Formula 1, as
described above, wherein

R₁ and R₃ are independently selected from the group consisting of:

- 1) H,
- 2) C₁₋₁₀ alkyl,
- 3) aryl, and
- 4) heterocyclyl,

5 said alkyl, aryl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

R₂ is:

- 1) H,
- 10 2) C₁₋₆ alkyl,
- 3) OH, or
- 4) halogen; and

R₁₀ is:

- 15 1) C₁₋₆ alkyl,
- 2) NR₇R₈,
- 3) O-C₁₋₆ alkyl,
- 4) aryl,
- 5) heterocyclyl,

20 said aryl and heterocyclyl is optionally substituted with from one to three members selected from R^a.

And yet another embodiment is the compound of Formula I wherein X is CH. A further embodiment is a compound as described in the second embodiment above wherein X is further defined as CH.

25 And still another embodiment is a compound as described in the second embodiment above wherein X is further defined as CH and R₁ is selected from the group consisting of:

- 1) H,
- 2) C₁₋₁₀ alkyl,

3) phenyl, and

4) heterocyclyl,

said alkyl, phenyl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

5

R₂ is H or C₁₋₆ alkyl;

R₃ is H or C₁₋₃ alkyl;

10 R₁₀ is:

1) C₁₋₆ alkyl,

2) NR₇R₈,

3) O-C₁₋₆ alkyl,

4) phenyl,

15

5) heterocyclyl,

said phenyl and heterocyclyl is optionally substituted with from one to three members selected from R^a; and

R^a is:

20

1) C₁₋₆ alkyl,

2) halogen,

3) NO₂,

4) OR,

5) NR₇R₈,

25

6) CN,

7) phenyl, or

8) heterocyclyl.

Yet another embodiment of the present invention is a compound which

- is 4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1-(3-piperidin-1-yl-propyl)-1H-pyridin-2-one;
- 1-(2-morpholin-4-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 5 1-(3-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 10 1-(2-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(1-dimethylamino-2-methyl-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 15 1-[2-(4-cyano-piperidin-1-yl-ethyl)]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-piperidin-1-yl-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-piperidin-1-yl-ethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 20 1-(2-morpholin-4-yl-ethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 25 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(2-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 30

- 1-(1-dimethylamino-2-methyl-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 5 1-[2-(4-cyano-piperidin-1-yl-ethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1-(3-piperidin-1-yl-propyl)-1H-pyrimidin-2-one;
- 1-(2-morpholin-4-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-10 2-one;
- 1-(3-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 15 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(2-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(1-dimethylamino-2-methyl-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-20 1H-pyrimidin-2-one; and
- 1-[2-(4-cyano-piperidin-1-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- or a pharmaceutically acceptable salt or stereoisomer thereof.

- Also included within the scope of the present invention is a
- 25 pharmaceutical composition which is comprised of a compound of Formula I as described above and a pharmaceutically acceptable carrier. The present invention also encompasses a method of treating or preventing cancer in a mammal in need of such treatment which is comprised of administering to said mammal a therapeutically effective amount of a compound of Formula I.
- 30 Preferred cancers for treatment are selected from cancers of the brain,

genitourinary tract, lymphatic system, stomach, larynx and lung. Another set of preferred forms of cancer are histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

Also included is a method of treating or preventing a disease in which angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula I. Such a disease in which angiogenesis is implicated is ocular diseases such as retinal vascularization, diabetic retinopathy, age-related macular degeneration, and the like.

Also included within the scope of the present invention is a method of treating or preventing inflammatory diseases which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula 1. Examples of such inflammatory diseases are rheumatoid arthritis, psoriasis, contact dermatitis, delayed hypersensitivity reactions, and the like.

Also included is a method of treating or preventing a tyrosine kinase-dependent disease or condition in a mammal which comprises administering to a mammalian patient in need of such treatment a therapeutically effective amount of a compound of Formula I. The therapeutic amount varies according to the specific disease and is discernable to the skilled artisan without undue experimentation.

A method of treating or preventing retinal vascularization which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Formula 1 is also encompassed by the present invention. Methods of treating or preventing ocular diseases, such as diabetic retinopathy and age-related macular degeneration, are also part of the invention. Also included within the scope of the present invention is a method of treating or preventing inflammatory diseases, such as rheumatoid arthritis, psoriasis, contact dermatitis and delayed

hypersensitivity reactions, as well as treatment or prevention of bone associated pathologies selected from osteosarcoma, osteoarthritis, and rickets.

The invention also contemplates the use of the instantly claimed compounds in combination with a second compound selected from the group

5 consisting of:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) retinoid receptor modulator,
- 4) a cytotoxic agent,
- 10 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor, and
- 15 10) another angiogenesis inhibitor.

Preferred angiogenesis inhibitors are selected from the group consisting of a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP inhibitor, an integrin blocker, interferon- α , interleukin-
20 12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, and an antibody to VEGF. Preferred estrogen receptor modulators are tamoxifen and raloxifene.

Also included in the scope of the claims is a method of treating
25 cancer which comprises administering a therapeutically effective amount of a compound of Formula 1 in combination with radiation therapy and/or in combination with a compound selected from the group consisting of:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 30 3) retinoid receptor modulator,

- 4) a cytotoxic agent,
5) an antiproliferative agent,
6) a prenyl-protein transferase inhibitor,
7) an HMG-CoA reductase inhibitor,
5 8) an HIV protease inhibitor,
9) a reverse transcriptase inhibitor, and
10) another angiogenesis inhibitor.

And yet another embodiment of the invention is a method of treating cancer which comprises administering a therapeutically effective
10 amount of a compound of Formula 1 in combination with paclitaxel or trastuzumab.

These and other aspects of the invention will be apparent from the teachings contained herein.

"Tyrosine kinase-dependent diseases or conditions" refers to
15 pathologic conditions that depend on the activity of one or more tyrosine kinases. Tyrosine kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion and migration, and differentiation. Diseases associated with tyrosine kinase activities include the proliferation of tumor cells, the pathologic neovascularization that
20 supports solid tumor growth, ocular neovascularization (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H.
25 Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers, with all possible isomers and mixtures thereof, including optical isomers, being included in the present invention. In addition, the compounds disclosed herein may exist as tautomers and both tautomeric
30 forms are intended to be encompassed by the scope of the invention, even

though only one tautomeric structure is depicted. For example, any claim to compound A below is understood to include tautomeric structure B, and vice versa, as well as mixtures thereof.



5

When any variable (e.g. aryl, heterocycle, R^1 , R^2 etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other occurrence. Also, combinations of substituents and variables are

10 permissible only if such combinations result in stable compounds.

It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily

15 available starting materials.

As used herein, "alkyl" is intended to include both branched, straight-chain, and cyclic saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C_1 - C_{10} , as in " C_1 - C_{10} alkyl" is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear, branched, or cyclic

20 arrangement. For example, " C_1 - C_{10} alkyl" specifically includes methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and so on, as well as cycloalkyls such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, tetrahydronaphthalene, methylenecyclohexyl, and so on. "Alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge.

25

If no number of carbon atoms is specified, the term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four

non-aromatic carbon-carbon double bonds may be present. Thus, "C₂-C₆ alkenyl" means an alkenyl radical having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl and cyclohexenyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Thus, "C₂-C₆ alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl and butynyl. As described above with respect to alkyl, the straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl, biphenyl, phenanthryl, anthryl or acenaphthyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring.

The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to: acridinyl, carbazolyl, cinnolinyl, quinoxaliny, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, indolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline. In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively.

As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro, fluoro, bromo and iodo. The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 5- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrathydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxaliny, tetrahydropyranyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidiny, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinoliny, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof.

The pharmaceutically acceptable salts of the compounds of this invention include the conventional non-toxic salts of the compounds of this invention as formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like: and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic,


glutamic, benzoic, salicylic, sulfanilic, 2-acetoxy-benzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

In certain instances, R^7 and R^8 are defined such that they can be taken together with the nitrogen to which they are attached to form a heterocyclic 5-10

5 membered saturated or unsaturated ring containing, in addition to the nitrogen atom, one to two additional heteroatoms selected from the group consisting of N, O and S, said ring optionally substituted with one or two substituents selected from R^a .

Examples of the 5-7 membered ring systems that can thus be formed include, but are not limited to the following:

10




The pharmaceutically acceptable salts of the compounds of this invention can be synthesized from the compounds of this invention which contain a

15 basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with

20 the appropriate inorganic or organic base.


The compounds of this invention may be prepared by employing reactions as shown in the following schemes, in addition to other standard manipulations that are known in the literature or exemplified in the experimental procedures. These schemes, therefore, are not limited by the compounds listed nor by
5 any particular substituents employed for illustrative purposes. Substituent numbering as shown in the schemes do not necessarily correlate to that used in the claims.

SCHEME 1



10





Generally, a method for the preparation of 3,6-diaryl pyrazolo(1,5-A)pyrimidines 3 comprises mixing a commercially available malondialdehyde compound 1 with a commercially available aminopyrazole 2 in an alcohol, such as
15 ethanol, methanol, isopropanol, butanol and the like, said alcohol containing catalytic quantities of an acid, such as acetic acid.

SCHEME 2

5 Scheme 2 illustrates one possible procedure for making 3,6-diaryl pyrazolo(1,5-A)pyrimidines **9** when the desired aminopyrazole is not commercially available. Compound **8** is obtained via the procedure described in Scheme 1 above. Subsequent treatment of **8** with a boronic acid derivative in the presence of a palladium catalyst provides the desired compound **9**.

10

SCHEME 3



5 Scheme 3 illustrates an alternative method for the preparation of 3,7 diarylpyrazolo(1,5-A)pyrimidines **21**. The commercially available ketone (**15**) and nitrile (**18**) are treated separately with dimethylformamidedi-methyl acetal (**16**) in

refluxing toluene to give produce 17 and 19, respectively. Compound 19 is then treated with hydrazine-hydrochloride in refluxing ethanol to yield the aminopyrazole 20. Compounds 17 and 20 are then treated with catalytic amounts of acetic acid in ethanol as described previously giving the desired of 3,7 diarylpyrazolo(1,5-
5 A)pyrimidines 21.

UTILITY

The instant compounds are useful as pharmaceutical agents for mammals, especially for humans, in the treatment of tyrosine kinase dependent
10 diseases. Such diseases include the proliferation of tumor cells, the pathologic neovascularization (or angiogenesis) that supports solid tumor growth, ocular neovascularization (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

The compounds of the instant invention may be administered to
15 patients for use in the treatment of cancer. The instant compounds inhibit tumor angiogenesis, thereby affecting the growth of tumors (J. Rak et al. *Cancer Research*, 55:4575-4580, 1995). The anti-angiogenesis properties of the instant compounds may also be useful in the treatment of certain forms of blindness related to retinal vascularization.

20 The disclosed compounds are also useful in the treatment of certain bone-related pathologies, such as osteosarcoma, osteoarthritis, and rickets, also known as oncogenic osteomalacia. (Hasegawa et al., *Skeletal Radiol.*, 28, pp.41-45, 1999; Gerber et al., *Nature Medicine*, Vol. 5, No. 6, pp.623-628, June 1999).

The compounds of this invention may be administered to mammals,
25 preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers or diluents, optionally with known adjuvants, such as alum, in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of
30 administration.

For oral use of a chemotherapeutic compound according to this invention, the selected compound may be administered, for example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and
5 lubricating agents, such as magnesium stearate, are commonly added. For oral administration in capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or
10 flavoring agents may be added. For intramuscular, intraperitoneal, subcutaneous and intravenous use, sterile solutions of the active ingredient are usually prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

The compounds of the instant invention may also be co-administered
15 with other well known therapeutic agents that are selected for their particular usefulness against the condition that is being treated. For example, the instant compounds may be useful in combination with known anti-cancer agents. Such known anti-cancer agents include the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents,
20 antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors.

"Estrogen receptor modulators" refers to compounds which interfere or inhibit the binding of estrogen to the receptor, regardless of mechanism. Examples of
25 estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyloxy)phenyl]-2H-1-benzopyran-3-yl)]-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenylhydrazone, and SH646.

"Androgen receptor modulators" refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism.

Examples of androgen receptor modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamideliazole, and abiraterone acetate.

- 5 "Retinoid receptor modulators" refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl)retinamide, N-4-carboxyphenyl retinamide,

- 10 "Cytotoxic agents" refer to compounds which cause cell death primarily by interfering directly with the cell's functioning or inhibit or interfere with cell myosis, including alkylating agents, tumor necrosis factors, intercalators, microtubulin inhibitors, and topoisomerase inhibitors.

- Examples of cytotoxic agents include, but are not limited to,
- 15 tirapazimine, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-
- 20 methylpyridine) platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum(II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-
- 25 deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755, and 4-demethoxy-3-deamino-3-aziridinyl-4-methylsulphonyl-daunorubicin.

- Examples of microtubulin inhibitors include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin,
- 30 dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476,

vinflunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl)benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, and BMS188797.

Some examples of topoisomerase inhibitors are topotecan,

- 5 hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrzolo[3,4,5-kl]acridine-2-(6H)propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]indolizino[1,2b]quinoline-10,13(9H,15H)dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100,
- 10 BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-
- 15 6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrzolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-
- 20 carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c]quinolin-7-one, and dimesna.


- "Antiproliferative agents" includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine,
- 25 trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-
 - 30 heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-

4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine, and 3-aminopyridine-2-carboxaldehyde thiosemicarbazone. "Antiproliferative agents" also includes monoclonal antibodies to growth factors, other than those listed under "angiogenesis inhibitors", such as trastuzumab.

"HMG-CoA reductase inhibitors" refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33. The terms "HMG-CoA reductase inhibitor" and "inhibitor of HMG-CoA reductase" have the same meaning when used herein.

Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see US Patent No. 4,231,938; 4,294,926; 4,319,039), simvastatin (ZOCOR®; see US Patent No. 4,444,784; 4,820,850; 4,916,239), pravastatin (PRAVACHOL®; see US Patent Nos. 4,346,227; 4,537,859; 4,410,629; 5,030,447 and 5,180,589), fluvastatin (LESCOL®; see US Patent Nos. 5,354,772; 4,911,165; 4,929,437; 5,189,164; 5,118,853; 5,290,946; 5,356,896), atorvastatin (LIPITOR®; see US Patent Nos. 5,273,995; 4,681,893; 5,489,691; 5,342,952) and cerivastatin (also known as rivastatin and BAYCHOL®; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase

inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention. An illustration of the lactone portion and its corresponding open-acid form is shown below as structures I and II.



In HMG-CoA reductase inhibitors where an open-acid form can exist, salt and ester forms may preferably be formed from the open-acid, and all such forms are included within the meaning of the term "HMG-CoA reductase inhibitor" as used herein.

- 10 Preferably, the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and most preferably simvastatin. Herein, the term "pharmaceutically acceptable salts" with respect to the HMG-CoA reductase inhibitor shall mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base, particularly those
- 15 formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc and tetramethylammonium, as well as those salts formed from amines such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, 1-p-chlorobenzyl-2-pyrrolidine-1'-yl-
- 20 methylbenzimidazole, diethylamine, piperazine, and tris(hydroxymethyl) aminomethane. Further examples of salt forms of HMG-CoA reductase inhibitors may include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate,
- 25 gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine,

hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, 5 tannate, tartrate, teoclate, tosylate, triethiodide, and valerate.

Ester derivatives of the described HMG-CoA reductase inhibitor compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy.

10 "Prenyl-protein transferase inhibitor" refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase). Examples of prenyl-protein transferase inhibiting compounds 15 include (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (-)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, 5(S)-n-butyl-1-(2,3-dimethylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone, 20 5(S)-n-Butyl-1-(2-methylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, 1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-2-piperazinone, 1-(2,2-diphenylethyl)-3-[N-(1-(4-cyanobenzyl)-1H-imidazol-5-ylethyl)carbamoyl]piperidine, 4-{5-[4-Hydroxymethyl-4-(4-chloropyridin-2-ylmethyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl}benzonitrile, 4-{5-[4-hydroxymethyl-4-(3-chlorobenzyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl}benzonitrile, 4-{3-[4-(2-oxo-2H-pyridin-1-yl)benzyl]-3H-imidazol-4-ylmethyl}benzonitrile, 4-{3-[4-(5-chloro-2-oxo-2H- 25 [1,2']bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl}benzonitrile, 4-{3-[4-(2-Oxo-

2H-[1,2']bipyridin-5'-ylmethyl}-3H-imidazol-4-ylmethyl}benzonitrile, 4-[3-(2-Oxo-1-phenyl-1,2-dihydropyridin-4-ylmethyl)-3H-imidazol-4-ylmethyl}benzonitrile, 18,19-dihydro-19-oxo-5*H*,17*H*-6,10:12,16-dimetheno-1*H*-imidazo[4,3-*c*][1,11,4]dioxazacyclo -nonadecine-9-carbonitrile, (±)-19,20-Dihydro-19-oxo-5*H*-
 5 18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxatriaza-cyclooctadecine-9-carbonitrile, 19,20-dihydro-19-oxo-5*H*,17*H*-18,21-ethano-6,10:12,16-dimetheno-22*H*-imidazo[3,4-*h*][1,8,11,14]oxatriazacycloeicosine-9-carbonitrile, and (±)-19,20-Dihydro-3-methyl-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-
 10 *k*][1,6,9,12]oxa-triazacyclooctadecine-9-carbonitrile.

Other examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U. S. Pat. No. 5,420,245, U. S. Pat. No. 5,523,430, U. S. Pat. No.
 15 5,532,359, U. S. Pat. No. 5,510,510, U. S. Pat. No. 5,589,485, U. S. Pat. No. 5,602,098, European Pat. Publ. 0 618 221, European Pat. Publ. 0 675 112, European Pat. Publ. 0 604 181, European Pat. Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Pat. No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO
 20 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Pat. No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO
 25 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U. S. Pat. No. 5,532,359. For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see European J. of Cancer, Vol. 35, No. 9, pp.1394-1401 (1999).

Examples of HIV protease inhibitors include amprenavir, abacavir, CGP-73547, CGP-61755, DMP-450, indinavir, nelfinavir, tipranavir, ritonavir, saquinavir, ABT-378, AG 1776, and BMS-232,632. Examples of reverse transcriptase inhibitors include delaviridine, efavirenz, GS-840, HB Y097,
 5 lamivudine, nevirapine, AZT, 3TC, ddC, and ddI.

"Angiogenesis inhibitors" refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR20),
 10 inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP inhibitors, integrin blockers, interferon- α , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (PNAS, Vol. 89, p. 7384 (1992); JNCI, Vol. 69, p. 475 (1982); Arch.
 15 Ophthalmol., Vol. 108, p.573 (1990); Anat. Rec., Vol. 238, p. 68 (1994); FEBS Letters, Vol. 372, p. 83 (1995); Clin. Orthop. Vol. 313, p. 76 (1995); J. Mol. Endocrinol., Vol. 16, p.107 (1996); Jpn. J. Pharmacol., Vol. 75, p. 105 (1997); Cancer Res., Vol. 57, p. 1625 (1997); Cell, Vol. 93, p. 705 (1998); Intl. J. Mol. Med., Vol. 2, p. 715 (1998); J. Biol. Chem., Vol. 274, p. 9116 (1999)), carboxyamidotriazole, combretastatin A-4,
 20 squalamine, 6-O-chloroacetyl-carbonyl-fumagillol, thalidomide, angiostatin, troponin-1, and antibodies to VEGF. (see, Nature Biotechnology, Vol. 17, pp.963-968 (October 1999); Kim et al., Nature, 362, 841-844 (1993)).

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-
 25 butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1H-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-
 pyrrolocarbonylimino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene
 30 disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_3$ integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the $\alpha_v\beta_5$ integrin, to compounds which

5 antagonize, inhibit or counteract binding of a physiological ligand to both the $\alpha_v\beta_3$ integrin and the $\alpha_v\beta_5$ integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins. The term also refers to antagonists of any combination of

10 $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_1\beta_1$, $\alpha_2\beta_1$, $\alpha_5\beta_1$, $\alpha_6\beta_1$ and $\alpha_6\beta_4$ integrins.

Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline,

15 N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-

20 dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

The instant compounds are also useful, alone or in combination with platelet fibrinogen receptor (GP IIb/IIIa) antagonists, such as tirofiban, to inhibit

25 metastasis of cancerous cells. Tumor cells can activate platelets largely via thrombin generation. This activation is associated with the release of VEGF. The release of VEGF enhances metastasis by increasing extravasation at points of adhesion to vascular endothelium (Amirkhosravi, *Platelets* 10, 285-292, 1999). Therefore, the present compounds can serve to inhibit metastasis, alone or in combination with GP

IIb/IIIa) antagonists. Examples of other fibrinogen receptor antagonists include abciximab, eptifibatide, sibrafiban, lamifiban, lotrafiban, cromofiban, and CT50352.

If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other
5 pharmaceutically active agent(s) within its approved dosage range. Compounds of the instant invention may alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a combination formulation is inappropriate.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the
10 compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

15 As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The term "therapeutically effective amount" as used herein means that
20 amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an
25 effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

The present invention also encompasses a pharmaceutical composition useful in the treatment of cancer, comprising the administration of a therapeutically effective amount of the compounds of this invention, with or without
30 pharmaceutically acceptable carriers or diluents. Suitable compositions of this

invention include aqueous solutions comprising compounds of this invention and pharmacologically acceptable carriers, e.g., saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's blood-stream by local bolus injection.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.

ASSAYS

The compounds of the instant invention described in the Examples were tested by the assays described below and were found to have kinase inhibitory activity. Other assays are known in the literature and could be readily performed by those of skill in the art. (see, for example, Dhanabal et al., *Cancer Res.* 59:189-197; Xin et al., *J. Biol. Chem.* 274:9116-9121; Sheu et al., *Anticancer Res.* 18:4435-4441; Ausprunk et al., *Dev. Biol.* 38:237-248; Gimbrone et al., *J. Natl. Cancer Inst.* 52:413-427; Nicosia et al., *In Vitro* 18:538-549).

VEGF RECEPTOR KINASE ASSAY

VEGF receptor kinase activity is measured by incorporation of radio-labeled phosphate into polyglutamic acid, tyrosine, 4:1 (pEY) substrate. The phosphorylated pEY product is trapped onto a filter membrane and the incorporation of radio-labeled phosphate quantified by scintillation counting.

MATERIALS

VEGF receptor kinase

The intracellular tyrosine kinase domains of human KDR (Terman, B.I. et al. Oncogene (1991) vol. 6, pp. 1677-1683.) and Flt-1 (Shibuya, M. et al. Oncogene (1990) vol. 5, pp. 519-524) were cloned as glutathione S-transferase (GST) gene fusion proteins. This was accomplished by cloning the cytoplasmic domain of the KDR kinase as an in frame fusion at the carboxy terminus of the GST gene. Soluble recombinant GST-kinase domain fusion proteins were expressed in *Spodoptera frugiperda* (Sf21) insect cells (Invitrogen) using a baculovirus expression vector (pAcG2T, Pharmingen).

Lysis buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.5% triton X-100, 10 % glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin and 1mM phenylmethylsulfonyl fluoride (all Sigma).

Wash buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.05% triton X-100, 10 % glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin and 1mM phenylmethylsulfonyl fluoride.

Dialysis buffer

50 mM Tris pH 7.4, 0.5 M NaCl, 5 mM DTT, 1 mM EDTA, 0.05% triton X-100, 50 % glycerol, 10 mg/mL of each leupeptin, pepstatin and aprotinin and 1mM phenylmethylsulfonyl fluoride.

10 X reaction buffer

200 mM Tris, pH 7.4, 1.0 M NaCl, 50 mM MnCl₂, 10 mM DTT and 5 mg/mL bovine serum albumin (Sigma).

30

Enzyme dilution buffer

50 mM Tris, pH 7.4, 0.1 M NaCl, 1 mM DTT, 10 % glycerol,
100 mg/mL BSA.

5 10 X Substrate

750 µg/mL poly (glutamic acid, tyrosine; 4:1) (Sigma).

Stop solution

30% trichloroacetic acid, 0.2 M sodium pyrophosphate (both
10 Fisher).

Wash solution

15% trichloroacetic acid, 0.2 M sodium pyrophosphate.

Filter plates

15 Millipore #MAFC NOB, GF/C glass fiber 96 well plate.

METHODA. Protein purification

1. Sf21 cells were infected with recombinant virus at a
20 multiplicity of infection of 5 virus particles/ cell and grown at 27 °C for 48
 hours.
2. All steps were performed at 4°C. Infected cells were
 harvested by centrifugation at 1000 X g and lysed at 4 °C for 30 minutes with
 1/10 volume of lysis buffer followed by centrifugation at 100,000Xg for 1 hour.
- 25 The supernatant was then passed over a glutathione Sepharose column
 (Pharmacia) equilibrated in lysis buffer and washed with 5 volumes of the same
 buffer followed by 5 volumes of wash buffer. Recombinant GST-KDR protein
 was eluted with wash buffer/10 mM reduced glutathione (Sigma) and dialyzed
 against dialysis buffer.

30

B. VEGF receptor kinase assay

1. Add 5 μ l of inhibitor or control to the assay in 50% DMSO.
2. Add 35 μ l of reaction mix containing 5 μ l of 10 X reaction buffer, 5 μ l 25 mM ATP/10 μ Ci [33 P]ATP (Amersham), and 5 μ l 10 X substrate.
3. Start the reaction by the addition of 10 μ l of KDR (25 nM) in enzyme dilution buffer.
4. Mix and incubate at room temperature for 15 minutes.
5. Stop by the addition of 50 μ l stop solution.
6. Incubate for 15 minutes at 4 $^{\circ}$ C.
7. Transfer a 90 μ l aliquot to filter plate.
8. Aspirate and wash 3 times with wash solution.
9. Add 30 μ l of scintillation cocktail, seal plate and count in a Wallac Microbeta scintillation counter.

Human Umbilical Vein Endothelial Cell Mitogenesis Assay

- Expression of VEGF receptors that mediate mitogenic responses to the growth factor is largely restricted to vascular endothelial cells. Human umbilical vein endothelial cells (HUVECs) in culture proliferate in response to VEGF treatment and can be used as an assay system to quantify the effects of KDR kinase inhibitors on VEGF stimulation. In the assay described, quiescent HUVEC monolayers are treated with vehicle or test compound 2 hours prior to addition of VEGF or basic fibroblast growth factor (bFGF). The mitogenic response to VEGF or bFGF is determined by measuring the incorporation of [3 H]thymidine into cellular DNA.

Materials**HUVECs**

HUVECs frozen as primary culture isolates are obtained from Clonetics Corp. Cells are maintained in Endothelial Growth Medium (EGM; Clonetics) and are used for mitogenic assays at passages 3-7.

5 Culture Plates

NUNCCLON 96-well polystyrene tissue culture plates (NUNC #167008).

Assay Medium

- 10 Dulbecco's modification of Eagle's medium containing 1 g/mL glucose (low-glucose DMEM; Mediatech) plus 10% (v/v) fetal bovine serum (Clonetics).

Test Compounds

- 15 Working stocks of test compounds are diluted serially in 100% dimethylsulfoxide (DMSO) to 400-fold greater than their desired final concentrations. Final dilutions to 1X concentration are made directly into Assay Medium immediately prior to addition to cells.

20 10X Growth factors

Solutions of human VEGF₁₆₅ (500 ng/mL; R&D Systems) and bFGF (10 ng/mL; R&D Systems) are prepared in Assay Medium.

10X [³H]Thymidine

- 25 [Methyl-³H]Thymidine (20 Ci/mmol; Dupont-NEN) is diluted to 80 uCi/mL in low-glucose DMEM.

Cell Wash Medium

Hank's balanced salt solution (Mediatech) containing 1 mg/mL bovine serum albumin (Boehringer-Mannheim).

5 Cell Lysis Solution

1 N NaOH, 2% (w/v) Na₂CO₃.

METHOD


1. HUVEC monolayers maintained in EGM are harvested by trypsinization and plated at a density of 4000 cells per 100 µL Assay Medium
- 10 per well in 96-well plates. Cells are growth-arrested for 24 hours at 37°C in a humidified atmosphere containing 5% CO₂.
2. Growth-arrest medium is replaced by 100 µL Assay Medium containing either vehicle (0.25% [v/v] DMSO) or the desired final concentration of test compound. All determinations are performed in triplicate.
- 15 Cells are then incubated at 37°C/5% CO₂ for 2 hours to allow test compounds to enter cells.
3. After the 2-hour pretreatment period, cells are stimulated by addition of 10 µL/well of either Assay Medium, 10X VEGF solution or 10X bFGF solution. Cells are then incubated at 37°C/5% CO₂.
- 20 4. After 24 hours in the presence of growth factors, 10X [³H]Thymidine (10 µL/well) is added.
5. Three days after addition of [³H]thymidine, medium is removed by aspiration, and cells are washed twice with Cell Wash Medium (400 µL/well followed by 200 µL/well). The washed, adherent cells are then solubilized by
- 25 addition of Cell Lysis Solution (100 µL/well) and warming to 37°C for 30 minutes. Cell lysates are transferred to 7-mL glass scintillation vials containing 150 µL of water. Scintillation cocktail (5 mL/vial) is added, and cell-associated radioactivity is determined by liquid scintillation spectroscopy.

Based upon the foregoing assays the compounds of formula I are inhibitors of VEGF and thus are useful for the inhibition of angiogenesis, such as in the treatment of ocular disease, e.g., diabetic retinopathy and in the treatment of cancers, e.g., solid tumors. The instant compounds inhibit VEGF-stimulated mitogenesis of human vascular endothelial cells in culture with IC₅₀ values between 0.01 - 5.0 μ M. These compounds also show selectivity over related tyrosine kinases (e.g., FGFR1 and the Src family; for relationship between Src kinases and VEGFR kinases, see Eliceiri et al., Molecular Cell, Vol. 4, pp.915-924, December 1999).

EXAMPLES

Examples provided are intended to assist in a further understanding of the invention. Particular materials employed, species and conditions are intended to be further illustrative of the invention and not limiting of the reasonable scope thereof.


EXAMPLE 1




3-(3-thiophenyl)-7-(4-pyridyl) pyrazolo(1,5-A)pyrimidine

A 13 x 100 mm reaction tube was charged with aminopyrazole (22) (16.5 mg, 0.100 mmol) dissolved in 0.500 mL EtOH and vinylogous amide (23) (17.6 mg, 0.100 mmol) dissolved in 0.200 mL EtOH. Glacial acetic acid (1 drop) was added and the reaction was heated to 80 °C for 14 h. An additional 0.100 mL of


glacial acetic acid was added and heating was continued for an additional 6 h. The sample was concentrated to dryness to provide the desired title compound. Analysis by mass spectrometry showed $[M+H]^+$ 279.2.



EXAMPLE 2




4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1-(3-piperidin-1-yl-propyl)-1H-pyridin-2-one


Step 1:

- 5 A solution of 2-(4-pyridyl)malondialdehyde (1.00 g, 6.70 mmol, 1 equiv) and 3-amino-4-phenylpyrazole (1.07 g, 6.72 mmol, 1.00 equiv) in absolute ethanol (50 mL) was heated at reflux for 2 h. The reaction mixture was allowed to cool to 23°C, and the resulting precipitate was filtered, washed with methanol (100 mL), and air dried to give 5 as a light yellow solid.
- 10 mp = 226-228°C. ¹H NMR (400 MHz, CDCl₃) δ 8.95 (d, 1H, *J* = 2.4 Hz), 8.85 (d, 1H, *J* = 2.4 Hz), 8.78 (atypical dd, 2H, *J* = 4.6, 1.6 Hz), 8.52 (s, 1H), 8.06 (dd, 2H, *J* = 8.0, 1.1 Hz), 7.55 (atypical dd, 2H, *J* = 4.6, 1.6 Hz), 7.48 (t, 2H, *J* = 7.9 Hz), 7.32 (td, 1H, *J* = 8.2, 0.9 Hz); anal. calcd for C₁₇H₁₂N₄: C, 74.98; H, 4.44; N, 20.58. Found C, 75.10; H, 4.56; N, 20.50.

15

Step 2:

- 5 A solution of 5 (300 mg, 1.10 mmol, 1 equiv) and 55% mCPBA (449 mg, 1.43 mmol, 1.30 equiv) in dichloromethane (50 mL) was stirred at 23°C for 5 h. The precipitate was filtered, washed with dichloromethane (50 ml) and air dried to give 6 as a bright yellow solid.
- ¹H NMR (400 MHz, (CD₃)₂SO) δ 9.73 (d, 1H, *J* = 2.3 Hz), 9.14 (d, 1H, *J* = 2.3 Hz), 8.86 (s, 1H), 8.37 (d, 2H, *J* = 7.0 Hz), 8.18 (d, 2H, *J* = 7.7 Hz), 8.01 (d, 2H, *J* = 7.3 Hz), 7.47 (t, 2H, *J* = 8.0 Hz), 7.28 (t, 1H, *J* = 7.4 Hz).
- 10

Step 3:


15

- A solution of 6 (70 mg, 0.24 mmol, 1 equiv) in acetic anhydride (10 mL) was heated at reflux for 16 h. The reaction mixture was allowed to cool to 23°C, then concentrated. To a solution of the residue in anhydrous methanol (15 mL) was added potassium carbonate (50 mg, 0.36 mmol, 1.5 equiv), and the resulting mixture was stirred at 23°C for 30 min. Excess potassium carbonate was removed by filtration, and the filtrate was concentrated. The residue was partitioned between water (100
- 20

mL) and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated to leave intermediate **7** as a brown solid.

¹H NMR (400 MHz, CDCl₃) δ 9.73 (d, 1H, *J* = 2.3 Hz), 9.14 (d, 1H, *J* = 2.3 Hz), 8.86 (s, 1H), 8.18 (d, 2H, *J* = 7.7 Hz), , 7.53 (d, 1H, *J* = 6.7 Hz), 7.47 (t, 2H, *J* = 7.8 Hz),
5 7.28 (t, 1H, *J* = 7.6 Hz), 6.92 (d, 1H, *J* = 1.2 Hz), 6.75 (dd, 1H, *J* = 6.7, 1.2 Hz).

Step 4:




10

A mixture of unpurified **7** (50 mg, 0.17 mmol, 1 equiv), 1-(3-chloropropyl)piperidine hydrochloride (41 mg, 0.21 mmol, 1.2 equiv), cesium carbonate (136 mg, 0.416 mmol, 2.4 equiv), and sodium iodide (31 mg, 0.21 mmol, 1.2 equiv) in DMF (5 mL) was heated at 50°C for 16 h. More 1-(3-chloropropyl)piperidine hydrochloride (34
15 mg, 0.17 mmol, 1.0 equiv), cesium carbonate (55 mg, 0.17 mmol, 1.0 equiv), and sodium iodide (25 mg, 0.17 mmol, 1.0 equiv) were added and the resulting mixture was heated at 60°C for 7 h. At this point, additional 1-(3-chloropropyl)piperidine

hydrochloride (34 mg, 0.17 mmol, 1.0 equiv), cesium carbonate (55 mg, 0.17 mmol, 1.0 equiv), and sodium iodide (25 mg, 0.17 mmol, 1.0 equiv) were added and heating (60°C) was continued for 48 h. The reaction mixture was then partitioned between water (50 mL) and ethyl acetate (3 x 50 mL). The combined organic layers were dried
5 over sodium sulfate and concentrated. The residue was purified by flash column chromatography (CHCl₃, sat'd with NH₃) to afford compound 8a as a yellow solid.
¹H NMR (400 MHz, CDCl₃) δ 8.88 (d, 1H, *J* = 2.3 Hz), 8.79 (d, 1H, *J* = 2.3 Hz), 8.51 (s, 1H), 8.05 (d, 2H, *J* = 7.7 Hz), , 7.56 (d, 1H, *J* = 7.0 Hz), 7.48 (t, 2H, *J* = 7.8 Hz), 7.31 (t, 1H, *J* = 7.4 Hz), 6.84 (d, 1H, *J* = 2.0 Hz), 6.39 (dd, 1H, *J* = 7.0, 2.1 Hz), 4.07
10 (t, 2H, *J* = 6.7 Hz), 2.36 (br m, 2H), 2.34 (t, 2H, *J* = 6.9 Hz), 2.00 (pentet, 2H, *J* = 6.8 Hz), 1.59 (brm, 6H), 1.45 (br m, 2H).

WHAT IS CLAIMED IS:

1. A compound in accordance with formula I:



I

or a pharmaceutically acceptable salt or stereoisomer thereof, wherein:

10 X is CH or N;

R₁ and R₃ are independently selected from the group consisting of:

- 1) H,
- 2) C₁₋₁₀ alkyl,
- 15 3) C₂₋₁₀ alkenyl,
- 4) C₂₋₁₀ alkynyl,
- 5) aryl,
- 6) halo,
- 7) OH, and
- 20 8) heterocyclyl,

said alkyl, alkenyl, alkynyl, aryl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

R₂ is:

- 5
- 1) H,
 - 2) C₁₋₆ alkyl,
 - 3) aryl,
 - 4) OH,
 - 5) NO₂,
 - 6) NH₂, or
 - 7) halogen;

10 R₅ is:

- 15
- 1) H,
 - 2) C₁₋₆ alkyl,
 - 3) OH,
 - 4) O-C₁₋₆ alkyl,
 - 5) halo,
 - 6) NH₂, or
 - 7) NO₂;

R₇ and R₈ are independently selected from the group consisting of:

- 20
- 1) H,
 - 2) C₁₋₁₀ alkyl,
 - 3) COR,
 - 4) COOR,
 - 5) aryl, and
 - 25 6) heterocyclyl,

said alkyl, aryl and heterocyclyl optionally substituted with R₉,
or NR₇R₈ are taken together to form a heterocyclic 5-10 membered
saturated or unsaturated ring containing, in addition to the nitrogen atom, one

to two additional heteroatoms selected from the group consisting of N, O and S, said ring optionally substituted with one or two substituents selected from R^a;

- 5 R₉ is aryl or heterocyclyl,
said aryl and heterocyclyl is optionally substituted with from one
to three members selected from R^a;

R₁₀ is:

- 10 1) H,
2) C₁₋₆ alkyl,
3) NR₇R₈,
4) O-C₁₋₆ alkyl,
5) aryl, or
6) heterocyclyl,

- 15 said alkyl, aryl, and heterocyclyl is optionally substituted with one to three
members selected from R^a;

R^a is:

- 20 1) C₁₋₁₀ alkyl,
2) halogen,
3) NO₂,
4) OR,
5) NR₇R₈,
6) CN,
25 7) aryl, or
8) heterocyclyl; and

R is H or C₁₋₆ alkyl.

2. A compound in accordance with Claim 1 wherein

R₁ and R₃ are independently selected from the group consisting of:

- 5
- 1) H,
 - 2) C₁₋₁₀ alkyl,
 - 3) aryl, and
 - 4) heterocyclyl,

said alkyl, aryl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

10

R₂ is:

- 1) H,
- 2) C₁₋₆ alkyl,
- 3) OH, or
- 15 4) halogen; and

R₁₀ is:

- 1) C₁₋₆ alkyl,
- 2) NR₇R₈,
- 20 3) O-C₁₋₆ alkyl,
- 4) aryl,
- 5) heterocyclyl,

said aryl and heterocyclyl is optionally substituted with from one to three members selected from R^a.

25

3. The compound of Claim 1 wherein X is CH.
4. The compound of Claim 2 wherein C is CH.

5. The compound of Claim 4 wherein

R₁ is selected from the group consisting of:

- 5
- 1) H,
 - 2) C₁₋₁₀ alkyl,
 - 3) phenyl, and
 - 4) heterocyclyl,

said alkyl, phenyl, and heterocyclyl is optionally substituted with one to three members selected from R^a;

10

R₂ is H or C₁₋₆ alkyl;

R₃ is H or C₁₋₃ alkyl;

15 R₁₀ is:

- 1) C₁₋₆ alkyl,
- 2) NR₇R₈,
- 3) O-C₁₋₆ alkyl,
- 4) phenyl,
- 20 5) heterocyclyl,

said phenyl and heterocyclyl is optionally substituted with from one to three members selected from R^a; and

R^a is:

- 25
- 1) C₁₋₆ alkyl,
 - 2) halogen,
 - 3) NO₂,
 - 4) OR,
 - 5) NR₇R₈,

- 6) CN,
- 7) phenyl, or
- 8) heterocyclyl.

- 5 6: A compound which is selected from the group consisting of:
- 4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1-(3-piperidin-1-yl-propyl)-1H-pyridin-2-one;
- 1-(2-morpholin-4-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 10 1-(3-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 15 1-(2-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(1-dimethylamino-2-methyl-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 20 1-[2-(4-cyano-piperidin-1-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-piperidin-1-yl-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-piperidin-1-yl-ethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 25 1-(2-morpholin-4-yl-ethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;

- 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 5 1-(2-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(1-dimethylamino-2-methyl-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 1-(3-dimethylamino-propyl)-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 10 1-[2-(4-cyano-piperidin-1-yl-ethyl)]-4-(3-thiophen-3-yl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyridin-2-one;
- 4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1-(3-piperidin-1-yl-propyl)-1H-pyrimidin-2-one;
- 15 1-(2-morpholin-4-yl-ethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(3-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(1-methyl-piperidin-3-ylmethyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 20 1-[3-(4-methylpiperazin-1-yl)-propyl]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 1-(2-dimethylamino-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- 25 1-(1-dimethylamino-2-methyl-propyl)-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one; and
- 1-[2-(4-cyano-piperidin-1-yl-ethyl)]-4-(3-phenyl-pyrazolo[1,5-a]pyrimidin-6-yl)-1H-pyrimidin-2-one;
- or a pharmaceutically acceptable salt or stereoisomer thereof.

7. A pharmaceutical composition which is comprised of a compound in accordance with Claim 1 and a pharmaceutically acceptable carrier.
- 5 8. A method of treating or preventing cancer in a mammal in need of such treatment which is comprised of administering to said mammal a therapeutically effective amount of a compound of Claim 1.
- 10 9. A method of treating cancer or preventing cancer in accordance with Claim 8 wherein the cancer is selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung.
- 15 10. A method of treating or preventing cancer in accordance with Claim 8 wherein the cancer is selected from histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.
- 20 11. A method of treating or preventing a disease in which angiogenesis is implicated, which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.
- 25 12. A method in accordance with Claim 11 wherein the disease is an ocular disease.
13. A method of treating or preventing retinal vascularization which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Claim 1.

14. A method of treating or preventing diabetic retinopathy which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of compound of Claim 1.
- 5 15. A method of treating or preventing age-related macular degeneration which is comprised of administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.
- 10 16. A method of treating or preventing inflammatory diseases which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1.
- 15 17. A method according to Claim 16 wherein the inflammatory disease is selected from rheumatoid arthritis, psoriasis, contact dermatitis and delayed hypersensitivity reactions.
- 20 18. A method of treating or preventing a tyrosine kinase-dependent disease or condition which comprises administering a therapeutically effective amount of a compound of Claim 1.
- 25 19. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier.
20. A process for making a pharmaceutical composition which comprises combining a compound of Claim 1 with a pharmaceutically acceptable carrier.
21. A method of treating or preventing bone associated pathologies selected from osteosarcoma, osteoarthritis, and rickets which

comprises administering a therapeutically effective amount of a compound of Claim 1.

22. The composition of Claim 7 further comprising a second
5 compound selected from the group consisting of:
- 1) an estrogen receptor modulator,
 - 2) an androgen receptor modulator,
 - 3) retinoid receptor modulator,
 - 4) a cytotoxic agent,
 - 10 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 7) an HMG-CoA reductase inhibitor,
 - 8) an HIV protease inhibitor,
 - 9) a reverse transcriptase inhibitor, and
 - 15 10) another angiogenesis inhibitor.

23. The composition of Claim 22, wherein the second
compound is another angiogenesis inhibitor selected from the group consisting
of a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor,
20 an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived
growth factor, an MMP inhibitor, an integrin blocker, interferon- α , interleukin-
12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole,
combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol,
thalidomide, angiostatin, troponin-1, and an antibody to VEGF.
25

24. The composition of Claim 22, wherein the second
compound is an estrogen receptor modulator selected from tamoxifen and
raloxifene.

25. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy.

5 26. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a compound selected from the group consisting of:

- 10
- 1) an estrogen receptor modulator,
 - 2) an androgen receptor modulator,
 - 3) retinoid receptor modulator,
 - 4) a cytotoxic agent,
 - 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 15 7) an HMG-CoA reductase inhibitor,
 - 8) an HIV protease inhibitor,
 - 9) a reverse transcriptase inhibitor, and
 - 10) another angiogenesis inhibitor.

20 27. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy and a compound selected from the group consisting of:

- 25
- 1) an estrogen receptor modulator,
 - 2) an androgen receptor modulator,
 - 3) retinoid receptor modulator,
 - 4) a cytotoxic agent,
 - 5) an antiproliferative agent,
 - 6) a prenyl-protein transferase inhibitor,
 - 30 7) an HMG-CoA reductase inhibitor,

- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor, and
- 10) another angiogenesis inhibitor.

5 28. A method of treating or preventing cancer which
comprises administering a therapeutically effective amount of a compound of
Claim 1 and paclitaxel or trastuzumab.

10 29. A method of treating or preventing cancer which
comprises administering a therapeutically effective amount of a compound of
Claim 1 and a GPIIb/IIIa antagonist.

30. The method of Claim 29 wherein the GPIIb/IIIa
antagonist is tirofiban.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/05903

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C07D 487/04; A61K 31/519
US CL :544/281; 514/258

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/281; 514/258

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAPLUS, CAOLD, BEILSTEIN, MARPAT

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94/29295 A1 (FUJISAWA PHARMACEUTICAL CO., LTD.) 22 December 1994, see entire document.	1-7
A	US 5,843,951 A (INOUE et al.) 01 December 1998, see entire document.	1-7
A, P	US 5,985,882 A (INOUE et al.) 16 November 1999, see entire document.	1-7
A	US 4,482,555 A (DORIA et al.) 13 November 1984, see entire document.	1-7

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 JUNE 2000

Date of mailing of the international search report

19 JUL 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MUKUND SHAH

Telephone No. (703) 308-1235